

# 2012 Childhood and Adolescent Immunization Clinics

## Medina County Health Department

Childhood and Adolescent  
Immunization Clinic  
4800 Ledgewood Drive, Medina  
Monday – Friday  
8:00 a.m. – 12:00 p.m.  
1:00 p.m. – 4:00 p.m.



### Spencer – Monthly

Spencer Town Hall  
112 N. Main Street  
2nd Wednesday  
1:00 p.m. – 4:00 p.m.

### Lodi – Monthly

Lodi Resource Center  
119 Bank Street  
3rd Wednesday  
1:00 p.m. – 4:00 p.m.



### Brunswick – Monthly

**New! Starting March 2012**  
Brunswick Recreation Center  
3637 Center Road  
1st Wednesday  
1:00 p.m. – 4:00 p.m.

### Wadsworth – Monthly

**Expanded Hours Begin February 2012**  
Summa Wadsworth Rittman  
Hospital Founder's Hall  
195 Wadsworth Road  
4th Wednesday  
New! 9:00 a.m. – 11:30 a.m.  
1:00 p.m. – 4:00 p.m.

- Appointments for immunizations are recommended. They can be made by calling 330-723-9688 or toll-free at 1-888-723-9688, option 1.
- A select number of adult vaccines are available by appointment.
- Women, Infants, and Children (WIC) is also available at the Wadsworth and Lodi immunization clinics. Appointments for WIC are required. They can be made at 330-723-9629 or toll-free at 1-888-723-9688, option 4.

#### Accepted Insurance:

- Medicaid
- Care Source
- Well Care
- Buckeye Community

#### Non-Medicaid vaccine administration fee:

- \$10 per vaccine for Medina County children and adolescents
- \$14.50 per vaccine for out-of-county children and adolescents

No child will be denied immunizations due to the inability to pay.  
Services are partially funded by your local property tax health levy.



4800 Ledgewood Drive, Medina, OH 44256  
330-723-9688 1-888-723-9688  
[www.medinahhealth.org](http://www.medinahhealth.org)

# When Do Children and Teens Need Vaccinations?

Age	HepB Hepatitis B	DTaP/Tdap Diphtheria, tetanus, pertussis (whooping cough)	Hib <i>Haemophilus influenzae</i> type b	IPV Polio	PCV Pneumococcal conjugate	RV Rotavirus	MMR Measles, mumps, rubella	Varicella Chickenpox	HepA Hepatitis A	HPV Human papillo- mavirus	MCV4 Meningococcal conjugate	Influenza Flu
Birth	✓											
2 months	✓ (1–2 mos)	✓	✓	✓	✓	✓						
4 months	✓	✓	✓	✓	✓	✓						
6 months	✓ (6–18 mos)	✓	✓	✓ (6–18 mos)	✓	✓	✓ (12–15 mos)	✓ (12–15 mos)	✓✓ (2 doses given 6 mos apart at age 12–23 mos)			✓  (One dose each fall or winter to all people ages 6 mos and older)
12 months		✓	✓		✓							
15 months		✓	✓		✓							
18 months		Catch-up	Catch-up		Catch-up							
19–23 months	Catch-up	Catch-up	Catch-up	Catch-up	Catch-up	Catch-up	Catch-up	Catch-up				
4–6 years		✓		✓		✓	✓					
7–10 years		Catch-up										
11–12 years		✓ Tdap			Catch-up			Catch-up	Catch-up	Catch-up	✓✓✓	✓
13–15 years		Catch-up (Tdap)								Catch-up		Catch-up
16–18 years										Catch-up		✓

**Please note:** Cases of pertussis (whooping cough) have increased in children, teens, and adults in the last few years. Tragically, some infants too young to be fully protected by vaccination have died. Ask your doctor or nurse if your children have received all the pertussis shots needed for his or her age. Also, if you haven't had your pertussis shot, you need to get one.

**What is “Catch-up?”** If your child’s vaccinations are overdue or missing, get your child vaccinated as soon as possible. If your child has not completed a series of vaccinations on time, he or she will need only the remainder of the vaccinations in the series. There’s no need to start over.